



Patient Name: _____

Date of Birth: _____

 PHYO
 CMC0027-001NS Rev. 8/2022

**Casimersen (AMONDYS 45)
 Infusion Therapy Plan**
Baseline Patient Demographic

To be completed by the ordering provider.

 Diagnosis: _____ Height: _____ cm Weight: _____ kg Body Surface Area: _____ (m²)

 NKDA - No Known Drug Allergies Allergies: _____
Therapy Plan orders extend over time (several visits) including recurring treatment.

Please specify the following regarding the entire course of therapy:

Duration of treatment: _____ weeks _____ months _____ unknown

 Treatment should begin: as soon as possible (within a week) within the month
****Plans must be reviewed / re-ordered at least annually. ******ORDERS TO BE COMPLETED FOR EACH THERAPY****ADMIT ORDERS** Height and weight Vital signs**HYPOTENSION DEFINED ADMIT** Nursing communication

Notify Provider for Hypotension:

Prior to starting infusion, please determine the patient's threshold for hypotension as defined by the following parameters. This information will be needed in the event of an infusion reaction.

Hypotension is defined as follows:

1 month to 1 year - systolic blood pressure (SBP) less than 70

1 year to 11 years - systolic blood pressure (SBP) less than 70 + (2 x age in years)

11 years to 17 years - systolic blood pressure (SBP) less than 90

OR any age - systolic blood pressure (SBP) drop of more than 30% from baseline.

Baseline systolic blood pressure (SBP) x 0.7 = value below defined as hypotension.

NURSING ORDERS

Please select all appropriate therapy

IV START NURSING ORDERS Insert Peripheral IV

Place PIV if needed or access IVAD if available.

 lidocaine 1% BUFFERED (J-TIP LIDOCAINE)

0.2 mL, INTRADERMAL, PRN

 when immediate procedure needed when procedure will take about 1 minute patient / family preference for procedure

 Administration Instructions: NOTE: Do not use this medication in patients with bleeding disorders, platelets \leq 20,000, or in patients taking anticoagulants, when accessing implanted ports or using a vein that will be utilized for chemotherapy administration, nor for pre-term infants or neonates.
 lidocaine - prilocaine (EMLA) cream

TOPICAL, PRN

 when more than 60 minutes are available before procedure when procedure will take more than 1 hour
 patient / family preference for procedure

Administration Instructions: NOTE: In children < 3 months of age, or < 5 kg in weight, maximum application time is 1 hour.

 lidocaine - tetracaine (SYNERA) patch

TOPICAL, PRN

 when 20 - 30 minutes are available before procedure when procedure will take more than 1 hour

 when anticipated pain is less than 5 mm from skin surface patient / family preference for procedure



Patient Name: _____

Date of Birth: _____

 PHYO
 CMC0027-001NS Rev. 8/2022

**Casimersen (AMONDYS 45)
 Infusion Therapy Plan**
ORDERS TO BE COMPLETED FOR EACH THERAPY
NURSING ORDERS, CONTINUED
 lidocaine with transparent dressing 4% kit

TOPICAL, PRN

 when 20 - 30 minutes are available before procedure when procedure will take more than 1 hour

 patient / family preference for procedure

 Heparin flush
heparin flush

10 - 50 units, INTRAVENOUS, PRN, IV line flush. Per protocol, heparin should not be used to flush peripheral IVs. This heparin flush should be used with all central lines including IVADs, with the exception of de-accessing the IVAD.

heparin flush

100 - 300 units, INTRAVENOUS, PRN, IV line flush. Per protocol, heparin should not be used to flush peripheral IVs. For use only when de-accessing IVADs.

 Sodium chloride flush
Sodium chloride flush 0.9% injection

1 - 20 mL, INTRAVENOUS, PRN, IV line flush

Sodium chloride - preservative free 0.9% injection

1 - 30 mL, INTRAVENOUS, PRN, IV line flush

PRE-PROCEDURE LABS
 Urinalysis

Unit collect

INTERVAL: Every 4 weeks

DURATION: Until Discontinued

 Cystatin C with CKiD eGFR

Unit collect

INTERVAL: Every 12 weeks

DURATION: Until Discontinued

 Protein Urine

Unit collect

INTERVAL: Every 12 weeks

DURATION: Until Discontinued

 Creatinine Urine

Unit collect

INTERVAL: Every 12 weeks

DURATION: Until Discontinued

PRE-MEDICATIONS
 Acetaminophen pre-medication 30 minutes prior (15 mg / kg, maximum 650 mg)
Nursing communication

Administer only one of the acetaminophen orders, suspension or tablets, do not give both.

acetaminophen suspension

15 mg / kg, ORAL, for 1 dose pre-medication, give 30 minutes prior to infusion

Dose: _____

acetaminophen tablet

15 mg / kg ORAL, for 1 dose pre-medication, give 30 minutes prior to infusion

Dose: _____



Patient Name: _____

Date of Birth: _____

PHYO
CMC0027-001NS Rev. 8/2022

Casimersen (AMONDYS 45) Infusion Therapy Plan

ORDERS TO BE COMPLETED FOR EACH THERAPY

PRE-MEDICATIONS

- Diphenhydramine pre-medication 30 minutes prior (1 mg / kg, maximum 50 mg)**

Nursing communication

Administer only one of the diphenhydrAMINE pre-medication orders, liquid, capsule or injection, do not give more than one of the orders as a pre-medication.

diphenhydrAMINE liquid

1 mg / kg, ORAL, for 1 dose pre-medication, give 30 minutes prior to infusion

Dose: _____

diphenhydrAMINE capsule

1 mg / kg ORAL, for 1 dose pre-medication, give 30 minutes prior to infusion

Dose: _____

diphenhydrAMINE injection

1 mg / kg, INTRAVENOUS, 1 dose pre-medication, give 30 minutes prior to infusion

Dose: _____

INTRA-PROCEDURE

- Vital Signs**

Baseline vitals prior to start of casimersen, then monitor vitals every 15 minutes during casimersen infusion and for 30 minutes after infusion completed. In the event of any hypersensitivity or other infusion related symptoms, the infusion should be stopped and notify the prescribing ordering provider.

- Physician Communication Order**

Recommended casimersen dose = 30 mg / kg. Please enter the dose of casimersen in 'mg' to facilitate prior authorization requirements. Vial size is 100 mg, if possible and clinically acceptable, round to nearest 100 mg.

- Casimersen in sodium chloride 0.9% 150 mL infusion**

INTRAVENOUS, at 150 mL / hr, for 1 dose. Infuse Casimersen over 60 minutes via in-line 0.2 micron filter. Do not mix or infuse with other medications. Refrigerate.

Dose: _____

- Therapy Appointment Request**

Please select department for the therapy appointment request:

Expires in 365 days

Dallas Special Procedures Plano Infusion Center Dallas Neurology

Please check casimersen for appropriate scheduling.

EMERGENCY MEDICATIONS

- Nursing communication**

1. Hives or cutaneous reaction only – no other system involvement

PATIENT IS HAVING A DRUG REACTION:

- a. Stop the infusion
- b. Give diphenhydramine as ordered
- c. Check heart rate, respiratory rate and blood pressure every 5 minutes until further orders from provider.
- d. Connect patient to monitor (cardiac / apnea, blood pressure and oxygen saturation) if not already on one
- e. Notify provider for further orders



Patient Name: _____

Date of Birth: _____

PHYO
CMC0027-001NS Rev. 8/2022

**Casimersen (AMONDYS 45)
Infusion Therapy Plan**

ORDERS TO BE COMPLETED FOR EACH THERAPY

EMERGENCY MEDICATIONS, CONTINUED

2. Hives or cutaneous reaction plus one other system, i.e. abdominal cramping, vomiting, hypotension, altered mental status, respiratory distress, mouth / tongue swelling

PATIENT IS HAVING ANAPHYLAXIS:

- a. Stop the infusion
- b. Call code – do not wait to give epinephrine
- c. Give epinephrine as ordered
- d. Notify provider
- e. Check heart rate, respiratory rate and blood pressure every 5 minutes until the code team arrives.
- f. Connect patient to monitor (cardiac / apnea, blood pressure and oxygen saturation), if not already on one.
- g. Give diphenhydramine once as needed for hives
- h. May repeat epinephrine every 5 minutes x 2 doses for persistent hypotension and respiratory distress with desaturation until code team arrives.
- i. May give albuterol as ordered for wheezing with oxygen saturation stable while waiting for code team, continue to monitor oxygen saturation.

Hypotension is Defined as Follows:

1 month to 1 year – systolic blood pressure (SBP) less than 70
 1 year to 11 years – systolic blood pressure (SBP) less than 70 + (2 x age in years)
 11 years to 17 years – systolic blood pressure (SBP) less than 90
 OR any age – systolic blood pressure (SBP) drop more than 30% from baseline.
 Baseline systolic blood pressure (SBP) x 0.7 = value below defined as hypotension.

**EPINEPHrine Injection Orderable For Therapy Plan
(AMPULE / EPI - PEN JR. / EPI - PEN) 0.01 mg / kg**

0.01 mg / kg, INTRAMUSCULAR, EVERY 5 MINUTES PRN, for anaphylaxis and may be repeated for persistent hypotension and respiratory distress with desaturation until the code team arrives, for 3 doses.

Use caution with PIV administration. This solution has a pH < 5, or a pH > 9, or an osmolality > 600 mOsm / L.

Dose: _____

**Cardio / Respiratory Monitoring
Rationale for Monitoring: High risk patient (please specify risk)**

- Clinically significant cardiac anomalies or dysrhythmias
- Recent acute life-threatening event
- Unexplained or acutely abnormal vital signs
- Artificial airway (stent, tracheostomy, oral airway)
- Acute, fluctuating or consistent oxygen requirements

Monitor Parameters (select all that apply): Heart rate Oxygen saturation Respiratory rate

Telemetry Required: Yes No

diphenhydrAMINE injection

1 mg / kg, INTRAVENOUS, ONCE PRN, for hives or cutaneous reaction, for 1 dose. Maximum dose = 50 mg per dose, 300 mg per day.

Dose: _____

Albuterol for aerosol

0.25 mg / kg., INHALATION ONCE PRN, for wheezing, but oxygen saturations stable while waiting for code team, continue to monitor oxygen saturation for 1 dose

Dose: _____

POST-PROCEDURE

Nursing communication

Flush PIV or IVAD with 20 mL 0.9% sodium chloride (250 mL bag) at the completion of the infusion.

Flush IVAD with saline and heparin flush per protocol prior to de-accessing IVAD.

Discontinue PIV prior to discharge.



Patient Name: _____

Date of Birth: _____

PHYO
CMC0027-001NS Rev. 8/2022

**Casimersen (AMONDYS 45)
Infusion Therapy Plan**

ORDERS TO BE COMPLETED FOR EACH THERAPY

POST-PROCEDURE, CONTINUED

Sodium chloride 0.9% infusion
INTRAVENOUS, at 0 - 25 mL / hour ONCE, for 1 dose.
Dose: _____

(circle one):
MD DO

Signature of Provider

Credentials

Date

Time

Printed Name of Provider