

STIMATE

Please read "Patient Guide to Therapy Control with STIMATE Nasal Spray" in its entirety.

Reasons Patients need Stimate:

- Heavy menstrual periods
- Bleeding from the mouth or nose
- Joint bleeding
- Bleeding in the muscle
- Preparation for procedures

Stimate Dosing:

-Your doctor will prescribe Stimate based on your child's weight.

Less than 110 pounds: One spray in **ONE** nostril.

Greater than 110 pounds: One spray in **BOTH** nostrils.

*STIMATE may only be used for 2 days in a row. If your child is still bleeding you will need to call 214-456-2379 and further instructions will be given.

-You must keep a record of the treatments you give your child. At the back of the pamphlet there is a place to record the dates of treatments given to your child.

Common Side Effects of this medication:

- Warm face/face turning slightly red
- Headaches
- Sleepiness

*Please decrease the amount of liquid your child drinks on the days he/she received STIMATE.

Your child may only have _____ ounces of liquid on the days he/she receives this medication.

Storage/Expiration:

- The bottle can be refrigerated or kept at room temperature. It should not be exposed to warmth or heat.
- You **MUST** throw away the bottle in 6 months after opening. Always write the date opened on the bottle.
- Discard bottle after 25 sprays. This does not include sprays to prime the bottle.

Priming Instructions:

-Before giving Stimate you will need to prime the bottle. Here are instructions on how to do that:

- 1) Remove the protective cap
- 2) Tilt the bottle so the lower end of the small tube is in the deepest part of the medicine.
- 3) Press down on the spray pump 4 times.

-The bottle only remains primed for 1 week. It must be primed if it has been more than 1 week since last use.

If your child does not respond to Stimate, your doctor will discuss other treatment options.

Please call 214-456-2379 for any questions or concerns regarding your child's bleeding disorder or this medication.