



Patient Name: _____

Date of Birth: _____

PHYO
CMC85049-002NS Rev. 11/2021

**CHST RITUXIMAB
(RITUXAN or BIOSIMILAR)
INFUSION THERAPY PLAN (NEPHROLOGY)**

Baseline Patient Demographic

To be completed by the ordering provider.

Diagnosis: _____ Height: _____ cm Weight: _____ kg Body Surface Area: _____ (m²)

NKDA - No Known Drug Allergies Allergies: _____

Therapy Plan orders extend over time (several visits) including recurring treatment.

Please specify the following regarding the entire course of therapy:

Duration of treatment: _____ weeks _____ months _____ unknown

Treatment should begin: as soon as possible (within a week) within the month

****Plans must be reviewed / re-ordered at least annually. ****

ORDERS TO BE COMPLETED FOR EACH THERAPY

ADMIT ORDERS

Height and weight

Vital signs

Hypotension Defined Admit

Nursing communication

Prior to starting infusion, please determine the patient's threshold for hypotension as defined by the following parameters. This information will be needed in the event of an infusion reaction occurring.

Hypotension is defined as follows:

1 month to 1 year - systolic blood pressure (SBP) less than 70

1 year to 11 years - systolic blood pressure (SBP) less than 70 + (2 x age in years)

11 years to 17 years - systolic blood pressure (SBP) less than 90

OR any age - systolic blood pressure (SBP) drop of more than 30% from baseline.

Baseline systolic blood pressure (SBP) x 0.7 = value below defined as hypotension.

PREGNANCY TESTS AT DALLAS AND PLANO

Nursing communication

Only one pregnancy test is necessary, based on facility capabilities. Please utilize the lab that is available per facility.

Patient requires a pregnancy test (based on organizational policy, female patients over 10 require a pregnancy test)

Pregnancy test, urine - POC

STAT, ONE TIME, for females > 10 years old. If positive, do NOT infuse and contact the ordering provider.

Gonodotropin chorionic (HCG) urine

STAT, ONE TIME, unit collect, for females > 10 years old. If positive, do NOT infuse and contact ordering provider.

NURSING ORDERS

Please select all appropriate therapy

IV START NURSING ORDERS

Insert peripheral IV / Access IVAD

Place PIV if needed or access IVAD if available

lidocaine 1% BUFFERED (J-TIP LIDOCAINE)

0.2 mL, INTRADERMAL, PRN

when immediate procedure needed when procedure will take about 1 minute patient / family preference for procedure

Administration Instructions: NOTE: Do not use this medication in patients with bleeding disorders, platelets ≤ 20,000, or in patients taking anticoagulants, when accessing implanted ports or using a vein that will be utilized for chemotherapy administration, nor for pre-term infants or neonates.



Patient Name: _____

Date of Birth: _____

PHYO
CMC85049-002NS Rev. 11/2021 **CHST RITUXIMAB
(RITUXAN or BIOSIMILAR)
INFUSION THERAPY PLAN (NEPHROLOGY)**

ORDERS TO BE COMPLETED FOR EACH THERAPY

NURSING ORDERS, CONTINUED

Please select all appropriate therapy

lidocaine - prilocaine (EMLA) cream

TOPICAL, PRN

when more than 60 minutes are available before procedure when procedure will take more than 1 hour

patient / family preference for procedure

Administration Instructions: NOTE: In children < 3 months of age, or < 5 kg in weight, maximum application time is 1 hour.

lidocaine - tetracaine (SYNERA) patch

TOPICAL, PRN

when 20 - 30 minutes are available before procedure when procedure will take more than 1 hour

when anticipated pain is less than 5 mm from skin surface patient / family preference for procedure

lidocaine with transparent dressing 4% kit

TOPICAL, PRN

when 20 - 30 minutes are available before procedure when procedure will take more than 1 hour

patient / family preference for procedure

Heparin flush

heparin flush

10 - 50 units, INTRAVENOUS, PRN, IV line flush. Per protocol, heparin should not be used to flush peripheral IVs. This heparin flush should be used with all central lines including IVADs, with the exception of de-accessing the IVAD.

heparin flush

100 - 300 units, INTRAVENOUS, PRN, IV line flush. Per protocol, heparin should not be used to flush peripheral IVs. For use only when de-accessing IVADs.

Sodium chloride flush

Sodium chloride flush 0.9% injection

1 - 20 mL, INTRAVENOUS, PRN, IV line flush

Sodium chloride - preservative free 0.9% injection

1 - 30 mL, INTRAVENOUS, PRN, IV line flush

PRE-PROCEDURE LABS

Complete Blood Count With Differential

STAT, ONE TIME, unit collect

DURATION: 4 treatments

PRE-MEDICATIONS

Acetaminophen pre-medication 30 minutes prior (15 mg / kg, maximum 650 mg)

Nursing communication

Administer only one of the acetaminophen orders, suspension or tablets, do not give both.

acetaminophen suspension

15 mg / kg, ORAL, for 1 dose pre-medication, give 30 minutes prior to infusion

Dose: _____

acetaminophen tablet

15 mg / kg ORAL, for 1 dose pre-medication, give 30 minutes prior to infusion

Dose: _____



Patient Name: _____

Date of Birth: _____

PHYO
CMC85049-002NS Rev. 11/2021

**CHST RITUXIMAB
(RITUXAN or BIOSIMILAR)
INFUSION THERAPY PLAN (NEPHROLOGY)**

ORDERS TO BE COMPLETED FOR EACH THERAPY

PRE-MEDICATIONS, CONTINUED

Diphenhydramine pre-medication 30 minutes prior (1 mg / kg, maximum 50 mg)

Nursing communication

Administer only one of the diphenhydrAMINE pre-medication orders, liquid, capsule or injection, do not give more than one of the orders as a pre-medication.

diphenhydrAMINE liquid

1 mg / kg, ORAL, for 1 dose pre-medication, give 30 minutes prior to infusion

Dose: _____

diphenhydrAMINE capsule

1 mg / kg ORAL, for 1 dose pre-medication, give 30 minutes prior to infusion

Dose: _____

diphenhydrAMINE injection

1 mg / kg, INTRAVENOUS, 1 dose pre-medication, give 30 minutes prior to infusion

Dose: _____

methyLPREDNISolone RTA infusion

2 mg / kg INTRAVENOUS, for 1 dose. Give 30 minutes prior to riTUXimAB Ready to administer by IV infusion. Doses > 15 mg / kg should be given over a minimum of 1 hour. (see protocol for monitoring parameters.)

Dose: _____

INTRA-PROCEDURE

Nursing communication

Adverse reactions may include fever, chills, rigors, hypotension and severe allergic reactions (anaphylaxis)

Vital signs

Obtain baseline vitals prior to start of riTUXimab infusion. Then monitor vitals 15 minutes after initiation of the infusion and 15 minutes after each rate change. Check vitals at the completion of the infusion and observe post infusion based on patient status.

Nursing communication

In the event of adverse reactions or other infusion related symptoms, the infusion should be stopped and the provider notified.

Initial infusion	First hour infusion rate	0.5 mg / kg / hr	Maximum rate: 50 mg / hr
	If no infusion related events observed after 60 minutes	Increase rate by 1 mg / kg / hr	Maximum rate: 100 mg / hr
	If no infusion related events observed after 30 minutes	Continue to increase rate by 0.5 mg / kg / hr every 30 minutes until maximum rate has been reached	Maximum rate: 400 mg / hr
Subsequent infusion	First hour infusion rate	1 mg / kg / hr	Maximum rate: 100 mg / hr
	If no infusion related events are observed after 30 minutes	Increase rate by 1 mg / kg / hr every 30 minutes until maximum rate has been reached	Maximum rate: 400 mg / hr

Physician communication order

Please round up riTUXimab dose, if clinically acceptable, to nearest 100 mg to minimize waste. Dose regimen of riTUXimab: 375 750 mg / m² every 2 weeks x 4 doses, (maximum dose 1,000 mg per treatment). Please enter the dose of riTUXimab in 'mg' to facilitate prior authorization requirements.

riTUXimab (RITUXAN or biosimilar) (Select one product below):

riTUXimab (RITUXAN) in sodium chloride 0.9% infusion
Intravenous, once, starting 0.5 hours after treatment start time
Final concentration should be 1 mg / mL

INTERVAL: 1 time a week

DURATION: 4 treatments

Dose: _____ mg

Sodium Chloride 0.9% **Dextrose** (Dilute to a final concentration of 1 - 4 mg / mL). **Dose:** _____ mL.

Initial infusion	First hour infusion rate	0.5 mg / kg / hr	Maximum rate: 50 mg / hr
	If no infusion related events observed after 60 minutes	Increase rate by 1 mg / kg / hr	Maximum rate: 100 mg / hr
	If no infusion related events observed after 30 minutes	Continue to increase rate by 0.5 mg / kg / hr every 30 minutes until maximum rate has been reached	Maximum rate: 400 mg / hr
Subsequent infusion	First hour infusion rate	1 mg / kg / hr	Maximum rate: 100 mg / hr
	If no infusion related events are observed after 30 minutes	Increase rate by 1 mg / kg / hr every 30 minutes until maximum rate has been reached	Maximum rate: 400 mg / hr



Patient Name: _____

Date of Birth: _____

PHYO

CMC85049-002NS Rev. 11/2021

**CHST RITUXIMAB
(RITUXAN or BIOSIMILAR)**
INFUSION THERAPY PLAN (NEPHROLOGY)
ORDERS TO BE COMPLETED FOR EACH THERAPY
NURSING ORDERS, CONTINUED
 ritiximab-abbs (TRUXIMA) in sodium chloride 0.9% infusion

INTERVAL: 1 time a week

DURATION: 4 treatments

 Intravenous, once, starting 0.5 hours after treatment start time
 Final concentration should be 1 mg / mL

Dose: _____ mg

 Sodium Chloride 0.9% Dextrose (Dilute to a final concentration of 1 - 4 mg / mL). Dose: _____ mL.

Initial infusion	First hour infusion rate	0.5 mg / kg / hr	Maximum rate: 50 mg / hr
	If no infusion related events observed after 60 minutes	Increase rate by 1 mg / kg / hr	Maximum rate: 100 mg / hr
	If no infusion related events observed after 30 minutes	Continue to increase rate by 0.5 mg / kg / hr every 30 minutes until maximum rate has been reached	Maximum rate: 400 mg / hr
Subsequent infusion	First hour infusion rate	1 mg / kg / hr	Maximum rate: 100 mg / hr
	If no infusion related events are observed after 30 minutes	Increase rate by 1 mg / kg / hr every 30 minutes until maximum rate has been reached	Maximum rate: 400 mg / hr

Therapy Appointment Request

Please select department for the therapy appointment request:

Expires in 365 days

 Dallas Special Procedures Plano Infusion Center Dallas Allergy Dallas Transplant Dallas Neurology

EMERGENCY MEDICATIONS
 Nursing communication

1. Hives or cutaneous reaction only – no other system involvement

PATIENT IS HAVING A DRUG REACTION:

- a. Stop the infusion
- b. Give diphenhydramine as ordered
- c. Check heart rate, respiratory rate and blood pressure every 5 minutes until further orders from provider.
- d. Connect patient to monitor (cardiac / apnea, blood pressure and oxygen saturation) if not already on one
- e. Notify provider for further orders

2. Hives or cutaneous reaction plus one other system, i.e. abdominal cramping, vomiting, hypotension, altered mental status, respiratory distress, mouth / tongue swelling

PATIENT IS HAVING ANAPHYLAXIS:

- a. Stop the infusion
- b. Call code – do not wait to give epinephrine
- c. Give epinephrine as ordered
- d. Notify provider
- e. Check heart rate, respiratory rate and blood pressure every 5 minutes until the code team arrives.
- f. Connect patient to monitor (cardiac / apnea, blood pressure and oxygen saturation), if not already on one.
- g. Give diphenhydramine once as needed for hives
- h. May repeat epinephrine every 5 minutes x 2 doses for persistent hypotension and respiratory distress with desaturation until code team arrives.
- i. May give albuterol as ordered for wheezing with oxygen saturation stable while waiting for code team, continue to monitor oxygen saturation.

Hypotension is defined as follows:

- 1 month to 1 year – systolic blood pressure (SBP) less than 70
- 1 year to 11 years – systolic blood pressure (SBP) less than 70 + (2 x age in years)
- 11 years to 17 years – systolic blood pressure (SBP) less than 90
- OR any age – systolic blood pressure (SBP) drop more than 30% from baseline.
- Baseline systolic blood pressure (SBP) x 0.7 = value below defined as hypotension.

 EPINEPHrine Injection Orderable For Therapy Plan
 (AMPULE / EPI - PEN JR. / EPI - PEN) 0.01 mg / kg

0.01 mg / kg, INTRAMUSCULAR, EVERY 5 MINUTES PRN, for anaphylaxis and may be repeated for persistent hypotension and respiratory distress with desaturation until the code team arrives, for 3 doses

Use caution with PIV administration. This solution has a pH < 5, or a pH > 9, or an osmolality > 600 mOsm / L.

Dose: _____



Patient Name: _____

Date of Birth: _____

PHYO
CMC85049-002NS Rev. 11/2021

**CHST RITUXIMAB
(RITUXAN or BIOSIMILAR)
INFUSION THERAPY PLAN (NEPHROLOGY)**

ORDERS TO BE COMPLETED FOR EACH THERAPY

EMERGENCY MEDICATIONS, CONTINUED

Cardio / Respiratory Monitoring
Rationale for Monitoring: High risk patient (please specify risk)

- Clinically significant cardiac anomalies or dysrhythmias
- Recent acute life-threatening event
- Unexplained or acutely abnormal vital signs
- Artificial airway (stent, tracheostomy, oral airway)
- Acute, fluctuating or consistent oxygen requirements

Monitor Parameters (select all that apply): Heart rate Oxygen saturation Respiratory rate
Telemetry Required: Yes No

diphenhydrAMINE injection

1 mg / kg, INTRAVENOUS, ONCE PRN, for hives or cutaneous reaction, for 1 dose. Maximum dose = 50 mg per dose, 300 mg per day.

Dose: _____

Albuterol for aerosol

0.25 mg / kg., INHALATION ONCE PRN, for wheezing, but oxygen saturations stable while waiting for code team, continue to monitor oxygen saturation for 1 dose

Dose: _____

POST - PROCEDURE

Nursing communication

Flush PIV or IVAD with 20 mL 0.9% sodium chloride (250 mL bag) at the completion of the infusion.
Flush IVAD with saline and heparin flush per protocol prior to de-accessing IVAD.
Discontinue PIV prior to discharge.

Sodium chloride 0.9% infusion

INTRAVENOUS at 0 - 25 mL / hr. ONCE, for 1 dose.

Dose: _____

(circle one):
MD DO

Signature of Provider

Credentials

Date

Time

Printed Name of Provider